



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Mangrove blue carbon stocks and dynamics are controlled by hydrogeomorphic settings and landuse change**

**Citation for published version:**

Sasmitho, SD, Sillanpää, M, Hayes, MA, Bachri, S, Saragisasmitho, MF, Sidik, F, Hanggara, B, Mofu, WY, Rumbiak, VI, Hendri, Taberima, S, Suhaemi, Nugroho, JD, Pattiasina, TF, Widagti, N, Barakalla, Rahajoe, JS, Hartantri, H, Nikijuluw, V, Jowey, RN, Heatubun, C, Zu Ermgassen, P, Worthington, TA, Howard, J, Lovelock, CE, Friess, DA, Hutley, LB & Murdiyarso, D 2020, 'Mangrove blue carbon stocks and dynamics are controlled by hydrogeomorphic settings and landuse change', *Global Change Biology*.  
<https://doi.org/10.1111/gcb.15056>

**Digital Object Identifier (DOI):**

[10.1111/gcb.15056](https://doi.org/10.1111/gcb.15056)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Global Change Biology

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



MR. SIGIT D. SASMITO (Orcid ID : 0000-0001-5864-8596)

DR. MATTHEW HAYES (Orcid ID : 0000-0002-2127-898X)

Article type : Primary Research Articles

## Mangrove blue carbon stocks and dynamics are controlled by hydrogeomorphic settings and land-use change

Sigit D. Sasmito<sup>1,2</sup>, Meriadec Sillanpää<sup>3,4</sup>, Matthew A. Hayes<sup>5</sup>, Samsul Bachri<sup>6</sup>, Meli F. Saragi-Sasmito<sup>2</sup>, Frida Sidik<sup>7</sup>, Bayu Hanggara<sup>2</sup>, Wolfram Y. Mofu<sup>8</sup>, Victor I. Rumbiak<sup>8</sup>, Hendri<sup>8</sup>, Sartji Taberima<sup>6</sup>, Suhaemi<sup>9</sup>, Julius D. Nugroho<sup>8</sup>, Thomas F. Pattiasina<sup>9</sup>, Nuryani Widagti<sup>7</sup>, Barakalla<sup>10,11</sup>, Joeni S. Rahajoe<sup>12</sup>, Heru Hartantri<sup>12</sup>, Victor Nikijuluw<sup>10</sup>, Rina N. Jowey<sup>8</sup>, Charlie Heatubun<sup>8</sup>, Philine zu Ermgassen<sup>13</sup>, Thomas A. Worthington<sup>14</sup>, Jennifer Howard<sup>15</sup>, Catherine E. Lovelock<sup>16</sup>, Daniel A. Friess<sup>3</sup>, Lindsay B. Hutley<sup>1</sup>, Daniel Murdiyarso<sup>2,17</sup>

<sup>1</sup>Research Institute for Environment and Livelihoods, Charles Darwin University, Casuarina 0810, Australia

<sup>2</sup>Center for International Forestry Research, Bogor 16115, Indonesia

<sup>3</sup>Department of Geography, National University of Singapore, 1 Arts Link, Singapore 117570, Singapore

<sup>4</sup>Green Forest Product and Tech. Pte. Ltd., Research Department, Singapore

<sup>5</sup>Australian Rivers Institute – Coast and Estuaries, Griffith University, Gold Coast, Australia

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/GCB.15056](https://doi.org/10.1111/GCB.15056)

This article is protected by copyright. All rights reserved

<sup>6</sup>Faculty of Agriculture, University of Papua, Manokwari, Indonesia

<sup>7</sup>Institute for Marine Research and Observation, Ministry of Marine Affairs and Fisheries, Bali, Indonesia

<sup>8</sup>Faculty of Forestry, University of Papua, Manokwari, Indonesia

<sup>9</sup>Faculty of Fisheries, University of Papua, Manokwari, Indonesia

<sup>10</sup>Conservation International Indonesia, Jakarta, Indonesia

<sup>11</sup>Present address at World Resources Institute Indonesia, Jakarta 12170, Indonesia

<sup>12</sup>Research Centre for Biology, Indonesian Institute of Sciences (LIPI), Cibinong, Indonesia

<sup>13</sup>Changing Oceans Group, School of Geosciences, University of Edinburgh, Edinburgh, UK

<sup>14</sup>Department of Zoology, University of Cambridge, Cambridge CB2 3QZ, UK

<sup>15</sup>Conservation International, USA

<sup>16</sup>School of Biological Sciences, The University of Queensland, Queensland, Australia

<sup>17</sup>Department of Geophysics and Meteorology, Bogor Agricultural University, Bogor, Indonesia

Corresponding author: Sigit D. Sasmito; email: sigitdeni.sasmito@cdu.edu.au; phone: +61 4 3409

5716

## ABSTRACT

Globally, carbon-rich mangrove forests are deforested and degraded due to land-use and land-cover change (LULCC). The impact of mangrove deforestation on carbon emissions has been reported on a global scale; however, uncertainty remains at subnational scales due to geographical variability and field data limitations. We present an assessment of blue carbon storage at five mangrove sites across West Papua Province, Indonesia, a region that supports 10% of the world's mangrove area. The sites are representative of contrasting hydrogeomorphic settings and also capture change over a 25 year of LULCC chronosequence. Field-based assessments were conducted across 255 plots covering undisturbed and LULCC-affected mangroves (0-, 5-, 10-, 15- and 25-year-old post-harvest or regenerating forests as well as 15-year-old aquaculture ponds). Undisturbed mangroves stored total ecosystem carbon stocks of 182–2730 (mean  $\pm$  SD:  $1087 \pm 584$ ) Mg C/ha, with the large variation driven by hydrogeomorphic settings. The highest carbon stocks were found in estuarine interior mangroves, followed by open coast interior, open coast fringe and estuarine interior forests. Forest harvesting did not significantly affect soil carbon stocks, despite an elevated dead wood density relative to undisturbed forests, but it did remove nearly all live biomass. Aquaculture conversion removed 60% of soil carbon stock and 85% of live biomass carbon stock, relative to reference sites. By contrast, mangroves left to regenerate for more than 25 years reached the same level of biomass carbon compared to undisturbed forests, with annual biomass accumulation rates of  $3.6 \pm 1.1$  Mg C/ha/year. This study shows that hydrogeomorphic setting controls natural dynamics of mangrove blue carbon stocks, while long-term land-use changes affect carbon loss and gain to a substantial degree. Therefore, current land-based climate policies must incorporate landscape and land-use characteristics, and their related carbon management consequences, for more effective emissions reduction targets and restoration outcomes.

**Keywords:** coastal wetlands, climate change mitigation, restoration, Paris Agreement, LULCC, Indonesia



## INTRODUCTION

Mangrove forests are one of the most productive and efficient long-term natural carbon sinks (Bouillon et al., 2008; Donato et al., 2011), and as such have been identified alongside seagrasses and saltmarshes as key 'blue carbon' ecosystems (Lovelock & Duarte, 2019). Mangroves have experienced large-scale deforestation and conversion to other land uses, particularly in Southeast Asia (Hamilton & Casey, 2016; Richards & Friess, 2016). Mangrove deforestation and conversion generates substantial carbon emissions (Atwood et al., 2017; Hamilton & Friess, 2018), accounting for a substantial proportion of greenhouse gas (GHG) emissions for some countries (Murdiyarso et al., 2015; Taillardat et al., 2018).

Recently, the conservation of mangrove carbon stocks has been promoted in global climate negotiations due to their potential contribution to mitigating GHG emissions. In response, the number of mangrove blue carbon assessments has increased rapidly over the past decade (Donato et al., 2011; Adame et al., 2013; Kauffman et al., 2014; Stringer et al., 2015; Nam et al., 2016; among many others). However, the majority of mangrove carbon studies have been conducted in natural or relatively undisturbed systems, making it difficult to generate estimates of carbon stock loss or recovery as a consequence of land-use change and restoration efforts (Sasmito et al., 2019). Estimates of carbon stock loss are further complicated by the fact that biomass and soil carbon vary substantially across climatic gradients (Simard et al., 2019) and geomorphological settings (Rovai et al., 2018; Twilley et al., 2018).

This study investigates the variation in mangrove carbon stocks across hydrogeomorphic settings, as well as their loss and recovery following land-use change in West Papua Province, Indonesia. We first assessed and compared total carbon stocks and other biophysical factors (i.e., forest structure and physicochemical soil properties) in undisturbed mangroves across different hydrogeomorphic settings. Second, we compared and identified the changes in carbon stocks between undisturbed mangrove forests and forests affected by land-use change (i.e., mangrove harvesting, regeneration and aquaculture) across a 25-year chronosequence. Third, we

calculated potential carbon stock loss and recovery resulting from these new land management practices. Our findings contribute to an enhanced understanding of current blue carbon stocks as well as the potential emissions and removals generated by mangrove management in an important region of the global mangrove cover, and can therefore be used to refine national carbon emissions calculations.

## **MATERIALS AND METHODS**

### **Hydrogeomorphic settings and land use in sampling sites**

The study was conducted in the Bintuni and Kaimana Regencies of West Papua Province, Indonesia. The Papua region, encompassing the provinces of West Papua and Papua, represents nearly 10% of global mangrove forest area (Hamilton & Casey, 2016), and Bintuni Bay in particular has been identified as a global hotspot for mangrove biomass (Simard et al., 2019). Generally, mangrove forests in Bintuni Bay are identified as a tidal estuarine hydrogeomorphic setting, with undisturbed mangrove stands reaching up to 30 m in height (Sillanpää et al., 2017; Simard et al., 2019). Nearly 30 true mangrove tree species have been recorded in Bintuni Bay (Kusmana & Onrizal, 2003).

Field sampling was carried out across five study sites (Table 1; Figure 1), which were divided into distinct hydrogeomorphic settings according to a macro-scale hydrogeomorphic typology developed by Worthington, et al. (2018) and overlaid with our sampling locations. The sampling sites were located in two mangrove hydrogeomorphic settings: estuarine and open coast (Figure 1). To understand spatial variation in biophysical and hydrological properties within the macro-scale typological units, a meso-scale typology, fringe and interior mangroves, was nested within the macro-scale typology. The meso-scale classifications were determined by distinct hydrodynamics and sediment supply characteristics between fringe and interior mangrove locations (Woodroffe et al., 2016). Subsequently, we compared carbon stocks, forest structure and soil properties of undisturbed mangrove across four hydrogeomorphic mangrove settings: estuarine fringe (EF), estuarine interior (EI), open coast fringe (OCF) and open coast interior (OCI) mangroves (see Table 1 for detailed descriptions of the sampling locations).

Undisturbed mangroves were located across all sampling sites (Table 1) and characterized by limited anthropogenic disturbance. In addition, three different land-use types, namely harvested mangrove, post-harvest or regenerating mangroves and aquaculture ponds at two sites (Bintuni Bay and Kaimana), were sampled for carbon stocks. In the post-harvest mangrove site in Bintuni Bay, mangrove stands were rotationally harvested for sustainable forestry products that more than 25 years old (Sillanpää et al., 2017). Forest stands at post-harvest mangrove sites were logged within the same year as we carried out our carbon stock field survey. Moreover, we assessed carbon stocks across different regenerating mangrove stands or rotational harvesting ages (5, 10, 15 and 25 years). We also performed a further carbon stock assessment within an aquaculture development in Kaimana, which was established in a former mangrove forest 15 years prior to our survey.

## **Field sampling and data analyses**

### *Sampling design*

We established 255 circular plots (each with an area of 314 m<sup>2</sup>, covering a total area of 8 ha) along 48 transects across the five study sites. The plots were distributed across four mangrove hydrogeomorphic settings (described in Table 1), with field assessments conducted between 2015 and 2018. Total ecosystem carbon stocks were assessed for four carbon stock pools: above-ground tree biomass carbon (AGBC), dead wood carbon, below-ground root biomass carbon (BGBC) and soil carbon. The sampling plot size and design were adapted from a globally applied protocol for mangrove ecosystem carbon stock assessment (Kauffman & Donato, 2012). A maximum of six circular plots were established along each transect perpendicular to the coastline or tidal creek. At each plot, we measured tree diameter, counted and measured dead wood diameter, as well as collected dead wood and soil samples.

### *Forest structure and biomass carbon stocks assessment*

We measured tree diameter at breast height (DBH), which was generally considered 130 cm above the forest floor or 30 cm above the highest prop root for *Rhizophora* spp. (Kauffman &

Donato, 2012). DBH was measured inside two different plots determined by DBH classes: circular plots of 10 m radius for trees with DBH > 5 cm and circular, nested plots of 2 m radius for mangrove saplings and seedlings with DBH < 5 cm. Standing dead tree status was documented following the dead tree definition of Kauffman & Donato (2012), and carbon stock correction factors were applied accordingly.

Tree and root biomasses were estimated from tree DBH data using species-specific allometric equations (Supplementary Information Table S1). Biomass was estimated by using standard carbon content factors of 47% and 39% for above-ground and below-ground biomass respectively, as described by Kauffman & Donato (2012), and was expressed as carbon stocks in Mg C/ha. We calculated stand basal area (m<sup>2</sup>/ha) by summing basal area (m<sup>2</sup>) for all trees across the surveyed area and dividing with plot area (ha). We estimated tree density (trees/ha) by counting tree quantities (trees) across the surveyed area and dividing by area (ha).

#### *Dead wood carbon pool*

We measured all dead, downed wood, including stem, branch and prop root debris lying on the forest floor, using the planar intercept technique described by Kauffman & Donato (2012). We classified dead wood into four classes based on its diameter (D): fine (D < 0.6 cm), small (0.6 cm < D < 2.5 cm), medium (2.5 cm < D < 7.5 cm) and large sound or rotten class (D > 7.5 cm). Two diagonal line transects were established and intersected in the midpoint of each circular plot. The DBH for large sound and rotten woody debris classes were measured, while all fine, small and medium classes were only recorded. Specific gravity of all woody debris classes was measured using the water displacement method, while carbon content was obtained from carbon and nitrogen (CN) elemental analysis. Means of quadratic mean diameter (cm), specific gravity (g/cm<sup>3</sup>) and carbon content (%) of the samples from our study are summarized in Supplementary Information Table S2. Woody debris carbon stocks of all classes were averaged from all diagonal line transects within the plot.

### *Soil carbon pool*

We collected soil samples from the center of each circular plot using a stainless steel Eijkelpamp peat soil auger. A 5 cm sample of sediment was extracted from the midpoint of fixed horizons at depths of 0–15, 15–30, 30–50, 50–100 and >100 cm along the core. Organic soil depths varied depending on hydrogeomorphic setting and degree of degradation. Specifically, in Bintuni Bay where estuarine mangroves occurred, typically with a deep soil organic matter layer, we extended soil sample collection up to 300 cm, and samples were extracted from the midpoint of 100–200 cm, and from the deepest layer. In total, we collected 1,068 soil samples from all study sites.

Soil samples were processed by oven drying at 60°C until constant weight was reached. Bulk density ( $\text{g}/\text{cm}^3$ ) was determined for each sample by dividing the dried weight (g) with the given soil auger volume ( $\text{cm}^3$ ). Samples were ground using a mortar and pestle and passed through a 0.5 mm sieve to remove large roots and inorganic debris. A CN elemental analysis was used to obtain carbon content of soil samples from Bintuni Bay, whereas a loss on ignition (LOI) approach was applied for samples collected from Arguni Bay, Buruway, Etna Bay and Kaimana.

Consequently, we corrected the carbon content of one-third of LOI-derived soil organic matter samples using CN elemental analysis data, and the correction factors were applied to the rest of LOI-derived soil organic matter. An inorganic carbon content correction was conducted by using the CN elemental analysis approach (Howard et al., 2014) and applied to one-fifth of total samples. Soil carbon stock ( $\text{Mg C}/\text{ha}$ ) was the final product of bulk density ( $\text{g}/\text{cm}^3$ ) multiplied with the corrected carbon content (%) scaled by depth intervals (cm). Carbon density ( $\text{mg C}/\text{cm}^3$ ) was calculated using data on bulk density and carbon content.

### *Carbon stock loss and recovery calculation*

We applied a Shapiro-Wilk normality and Levene's homogeneity test prior to statistical comparisons, with a logarithmic and square root transformation applied if data were not normally distributed and failed the homogeneity test. When the data were normally distributed,

we used a one-way analysis of variance (ANOVA) and Bonferroni multiple comparison test to compare carbon stocks, forest structure and soil properties between sampling plots and LULCC types. We applied a nonparametric Kruskal Wallis and Wilcoxon rank sum multiple comparison test for datasets that were not distributed normally and homogeneously. When carbon stocks at each pool were statistically different ( $p < 0.05$ ), a carbon stocks difference approach was applied to estimate carbon stock loss and recovery impacted by land uses.

Carbon stock loss and recovery were estimated by subtraction of carbon stock pools between undisturbed reference mangrove forest and land-use affected sites (Kauffman et al., 2017; Kauffman et al., 2018; Arifanti et al., 2019). The reference forest was the one closest to each land-use type, within the same study site. Reference forests for rotational harvesting and regeneration land-use types were in Bintuni Bay. In addition, reference forest for aquaculture was chosen from Kaimana. For the soil carbon pool assessment, we standardized soil carbon stocks using the soil mass equivalent approach modified from Ellert et al. (2007). The minimum soil mass value at Bintuni and Kaimana, respectively, was used to standardize soil carbon stocks for each soil layer. This standardization allowed the reduction of uncertainty sourced from soil compaction impacted by land-use change.

To determine the best variables to predict total ecosystem carbon stocks, we used simple and multiple generalized linear regression models (R function 'glm') and applied at a plot level dataset (*sensu* Paz et al., 2016). We first defined *a priori* variables which may control total ecosystem carbon stocks, namely hydrogeomorphic setting, land-use change, basal area, tree density, number of tree species, bulk density, carbon content, carbon density, and soil depth. We used the R function 'dredge' to select the best regression models based on their AICc (corrected Akaike Information Criterion). The best fit models that were considered in the results were models with  $\Delta AICc < 4$ . All raw data from this study are accessible through the CIFOR Dataverse digital repository (Sasmito et al., 2019b, 2019c, 2019d), and all R code used for statistical analysis can be found in the online GitHub repository (<https://github.com/ssasmito/Papua-mangrove-blue-carbon>).

## RESULTS

### Carbon stocks variation across undisturbed mangroves

Total carbon stocks of undisturbed mangroves in Papua ranged between 182–2730 (mean  $\pm$  SD here after:  $1087 \pm 584$ ) Mg C/ha (Figure 2a). Total carbon stocks were significantly different among sites ( $p < 0.05$ ) (Figure 2a) and hydrogeomorphic settings ( $p < 0.05$ ) (Figure 2b). Across sampling sites, the largest mean of total carbon stocks was obtained at Arguni Bay ( $1686 \pm 564$  Mg C/ha), while the lowest was found at Kaimana ( $645 \pm 418$  Mg C/ha). In relation to the coastal hydrogeomorphic setting (Figure 2b), estuarine interior mangroves stored the largest total carbon stocks ( $1480 \pm 552$  Mg C/ha). By contrast, estuarine fringe mangrove settings stored the lowest total carbon stocks, with  $432 \pm 193$  Mg C/ha or only one-third of the carbon stocks found in estuarine interior settings. In open coast mangroves, total carbon stocks between fringe and interior settings were similar, at  $865 \pm 72$  and  $867 \pm 216$  Mg C/ha ( $p > 0.05$ ).

Between carbon pools, 89% of total carbon stocks were stored in the soil, while 10% and 1% were in biomass and dead wood, respectively. Both the total soil carbon stocks and stocks in the top 100 cm differed significantly among sites and hydrogeomorphic settings ( $p < 0.05$ ) (Figure 2a–2b, Supplementary Information Table S3). Total soil carbon stocks in Arguni and Bintuni Bay (estuarine interior mangroves) were twice as large as stock estimates at Buruway, Etna Bay and Kaimana (open coast and estuarine fringe mangroves). This pattern was associated with deeper soil organic profiles (up to 300 cm) in estuarine interior mangroves compared to the other hydrogeomorphic settings in this study (Table 1). In estuarine mangroves, 62% of the total soil carbon stocks were distributed in deeper soil layers ( $> 100$  cm) rather than the upper layer (top 100 cm). By contrast, deeper soil layers across other mangrove settings only contributed 39% of the total soil carbon stock.

The overall mean of above- and below-ground live biomass carbon stocks (AGBC and BGBC) were  $96 \pm 65$  and  $17 \pm 16$  Mg C/ha, respectively, with relatively larger stocks observed in Bintuni Bay,

Buruway and Etna compared to Kaimana ( $p < 0.05$ ). Across hydrogeomorphic settings, AGBC was significantly different ( $p < 0.05$ ), while BGBC was similar ( $p = 0.41$ ) (Supplementary Information Table S3). We also observed the largest dead wood carbon stocks in Bintuni Bay ( $20 \pm 14$  Mg C/ha) and the lowest in Kaimana ( $2 \pm 3$  Mg C/ha) ( $p < 0.05$ ), however, significant variation between hydrogeomorphic settings was not observed ( $p = 0.53$ ) (Supplementary Information Table S3).

Mangrove forest structural variables, such as basal area and tree density, varied significantly across undisturbed mangrove sites ( $p < 0.05$ ) (Figure 3a and Figure 3c). Mean basal area in Bintuni Bay, Buruway and Etna Bay was greater than in Arguni and Kaimana ( $p < 0.05$ ) (Figure 3a). The highest mean ( $34 \pm 13$  m<sup>2</sup>/ha) tree density was observed in Bintuni Bay, while the lowest ( $13 \pm 11$  m<sup>2</sup>/ha) was in Kaimana ( $p < 0.05$ ) (Figure 3c). However, across hydrogeomorphic settings, basal area and tree density were not significantly different ( $p > 0.05$ ) (Figure 3b and Figure 3d).

In addition, soil carbon density also differed significantly between sampling sites ( $p < 0.05$ ) (Figure 3e), but was similar across hydrogeomorphic settings ( $p > 0.05$ ) (Figure 3f). The largest mean soil carbon density was found in Bintuni Bay with  $46 \pm 10$  g C/cm<sup>3</sup>. Soil bulk density mean varied significantly between sampling sites and hydrogeomorphic settings ( $p < 0.05$ ) (Figure 3g and Figure 3h). Kaimana had the largest mean of soil bulk density ( $0.55 \pm 0.14$  g/cm<sup>3</sup>), while the lowest was observed in Arguni Bay ( $0.35 \pm 0.15$  g/cm<sup>3</sup>). Across hydrogeomorphic settings, estuarine fringe mangrove had the greatest mean of soil bulk density ( $0.56 \pm 0.24$  g/cm<sup>3</sup>), and the lowest was found in the estuarine interior mangrove setting ( $0.42 \pm 0.19$  g/cm<sup>3</sup>). Similar to bulk density, soil carbon content was significantly different between sampling sites and hydrogeomorphic settings ( $p < 0.05$ ) (Figure 3i and 3j). The largest carbon content was observed in Arguni Bay ( $19\% \pm 11\%$ ) and estuarine interior mangrove setting ( $15\% \pm 10\%$ ). Further, soil bulk and carbon densities increased significantly with soil depth ( $p < 0.05$ ) (Figure 4a and Figure 4c). By contrast, soil carbon content was similar across depths ( $p > 0.05$ ) (Figure 4b). However, within the extended soil sample collection in Bintuni Bay (>100 cm depth), at approximately 150 and



300 cm, all soil properties (bulk density, carbon content and carbon density) were significantly different ( $p < 0.05$ ) (Supplementary Information Table S3).

The best regression model identified in the generalized linear model analysis suggested that soil depth was the most important variable to predict variation in total ecosystem carbon stocks (indicated by the lowest  $p$ -value), followed by stand basal area, soil carbon density and bulk density (Supplementary Table S4). The ten best multiple regression models (out of 446 possible models) indicate hydrogeomorphic setting, land-use change, soil depth, stand basal area, and soil carbon density to be the optimal variables to describe variation in total ecosystem carbon stocks (Supplementary Table S4). These variables were selected by all of ten best models, while other variables such as soil bulk density was only selected by six models, soil carbon content by four models, and tree density as well as number of mangrove species by three models, respectively.

### **Carbon stock loss and recovery following land-use changes**

Rotational harvesting of mangrove forest in Bintuni Bay resulted in a nearly complete loss of live biomass carbon stocks and a 99% increase of dead wood (Figure 5a). The mean of dead wood carbon stocks at a 0-year-old post-harvest forest site was  $40 \pm 15$  Mg C/ha, compared to only  $20 \pm 14$  Mg C/ha at undisturbed reference forests (Figure 5a). Overall, rotational forest harvesting generated 75% net carbon stock losses due to combined stock changes of live biomass and dead wood carbon pools (Figure 5b). By contrast, there was little significant change of soil carbon stocks resulting from rotational harvesting, particularly between 0–200 cm upper soil layers ( $p > 0.05$ ) (Figure 5a and Figure 5b). Nevertheless, soil carbon stocks at the deepest soil layer (>200 cm) between these two land uses were significantly different ( $p < 0.05$ ) (Figure 5b), in which we observed larger carbon stocks in undisturbed than in 0-year-old post-harvest forests.

Mangrove regeneration over 25 years resulted in a mean recovery of biomass carbon stocks of  $3.6 \pm 1.1$  Mg C/ha/year. Compared to reference sites, live biomass carbon stocks at 25-year-old regenerated forests were not significantly different ( $p > 0.05$ ) (Figure 5b). Across forests that had

regenerated for 10 and 25 years, dead wood carbon stocks were lower than in reference forests ( $p < 0.05$ ) (Figure 5b). Carbon stocks at 0–200 cm upper soil layers across all regenerated forests were similar compared to reference stands ( $p > 0.05$ ) (Figure 5a and Figure 5b).

In Kaimana, live biomass and soil carbon stocks were significantly different between sites converted to aquaculture and undisturbed reference sites ( $p < 0.05$ ) (Figure 5a and Figure 5b). Mangrove conversion to aquaculture resulted in live biomass carbon stocks losses of 85% (Figure 5b). In addition, aquaculture conversion generated a soil carbon stocks decrease of ~60% at 0–50 cm soil layers (Figure 5b). Combining carbon stock losses from live biomass and soil carbon pools, mangrove to aquaculture conversion reduces carbon stocks by 66%.

## DISCUSSION

### Hydrogeomorphic settings control undisturbed mangrove carbon stocks

Across undisturbed mangroves in West Papua, Indonesia, variation in total ecosystem carbon stocks is strongly influenced by the underlying hydrogeomorphic setting. For example, in Bintuni and Arguni Bays, where mangroves are characterized by a tidally dominated estuarine setting, total carbon stocks were larger compared to those located in open coast settings, such as in Etna Bay (Figure 2). An estuarine mangrove setting is typically supported by extensive allochthonous sediment supply and resulting accommodation space, creating the spatial and vertical room within which mangrove sediments may accumulate (Woodroffe et al., 2016), leading to the maintenance of soil carbon burial rates over historical millennia timescales (Rogers et al., 2019). This pattern, however, may not be observed in open coast settings, typically characterized by a lower tidal range, steeper coastal profile and thus less accommodation space (Woodroffe et al., 2016). At the meso-scale, distinct hydrodynamic differences (tidal inundation and flushing) between fringe and interior forest locations also generated substantial carbon stocks variation, specifically within a large macro-tidal estuarine setting, such as in Bintuni Bay. Fringe mangrove locations are prone to seasonally driven sediment dynamics, while interior locations experience less tidal flushing, allowing persistent rates of organic-rich sedimentation and increased

accumulated autochthonous carbon inputs (Krauss et al., 2014; Sasmito et al., 2020). As a result, estuarine interior mangroves stored the largest ecosystem carbon stocks compared to the other hydrogeomorphic settings observed in this study (Figure 2b).

Similar to previous mangrove carbon stock assessments (Donato *et al.*, 2011; Murdiyarso *et al.*, 2015), the soil carbon pool accounted for the majority (89%) of carbon, relative to the other carbon pools, across mangrove settings (Figure 2). For instance, total carbon stocks in estuarine interior mangrove sites consisted of 94% soil carbon, while across the rest of the mangrove settings, the soil carbon pool represented between 71% and 85%. The larger proportion of the soil carbon pool in estuarine interior mangroves is attributed to the higher carbon content (Figure 3j) and deeper organic soil layers (Table 1), a function of the complex interaction between sediment supply, accommodation space, hydrodynamics and biomass productivity. Overall, our findings suggest that the hydrogeomorphic setting substantially controls soil carbon and therefore, total carbon stocks in mangrove ecosystems.

#### **Effect of soil properties and forest structure on carbon stocks**

Comparison of the outcomes of multiple regression analyses suggest that total ecosystem carbon stocks are strongly influenced by variation in soil properties (soil depth, bulk density and carbon density), forest structure (basal area), hydrogeomorphic settings, and land-use change. Soil depth was the most significant variable explaining variation in total ecosystem carbon stocks, suggesting that deeper soil depth layers have the larger carbon stocks. These findings are consistent with recent global scale assessments (Kauffman et al., 2020), in which carbon stock estimates could be double those currently reported if soil carbon stocks > 100 cm are included in the blue carbon stock estimates. The results of our analyses also clarify that both hydrogeomorphic setting and land use change are equally significant factors in contributing to variation in total ecosystem carbon stocks.

Soil physicochemical properties, including carbon density, soil bulk density and carbon content, are the important factors contributing to the variation in soil carbon stocks. Previous assessments of mangrove soil carbon stocks typically extrapolate the size of the soil pool to 300 cm (Donato et al., 2011; Kauffman et al., 2011; Adame et al., 2018), rather than use measured observations at these depths. Extrapolation of soil carbon to deeper soil depths is usually based on assessments of bulk density and carbon content from ~75 cm depth (midpoint between 50–100 cm). However, with collection of soil samples to 300 cm, as we achieved at Bintuni Bay, we found that carbon content in the deeper soil layers (>100 cm) was two to three times lower than in upper layers (<100 cm) (Supplementary Information Table S3). This pattern was also observed for bulk density, in which bulk density at ~300 cm was nearly double of that observed in 0–50 cm layers. Clearly, collection of samples from deeper soil layers enhances the precision of stock estimates and the understanding of soil carbon variability with depth. Despite high costs, high density replicated sampling at depth and across settings must be incorporated into carbon stock assessments. This will reduce uncertainties associated with variations in soil carbon over hydrogeomorphic settings and with future land-use changes.

In addition, AGBC between mangrove sites and settings was also significantly different (Supplementary Information Table S3) despite tree density and basal area being similar among settings (Figure 3b and 3d). The variation of AGBC may be attributed to variation in species composition and the application of species-specific tree biomass allometric equations (Supplementary Information Table S1). In contrast, BGBC was similar between mangrove settings (Supplementary Information Table S3). In this case, species specific equations were not fully applied because suitable species specific equations from relevant geographical and climatic conditions were not available, specifically for multiple species including *Sonneratia* spp. and *Xylocarpus* spp. (Supplementary Information Table S1). The variability in the dominant mangrove species among sites and settings results in variation in biomass carbon stocks. Our findings also suggest that future refinement of mangrove carbon stocks assessments requires improved tree allometric equations, particularly for the belowground root carbon pool (Adame et al., 2017).

## **Mangrove harvesting and aquaculture conversion generates carbon stock losses**

Our findings suggest that land management practices such as forest harvesting and small-scale aquaculture reduce carbon stocks substantially, with the degree of reduction dependent on the type of land-use change. This is similar to the broad conclusions derived from a recent global-scale systematic review and meta-analysis (Sasmito, et al., 2019) but extends this further by region specific emissions factors. Within Bintuni Bay harvested forests, live vegetation biomass carbon stocks were nearly zero due to timber extraction. Nevertheless, not all vegetation was extracted successfully, leaving some dead stumps behind, and consequently there was a 100% increase of dead wood material left on the ground (Figure 5a). Our assessment was unable to quantify the dead biomass carbon located in the below-ground roots of logged stands, suggesting that further quantification of this carbon pool may improve current understanding of carbon stock loss associated with forest harvesting.

There was no statistical difference in soil carbon stocks of upper soil layers (0–200 cm) between logged forests and undisturbed reference sites, suggesting that most soil carbon stocks remain preserved due to minimal soil disturbance during the harvesting process. However, logged forest areas are continuously emitting GHGs as a result of decomposition of below-ground and dead biomass, as observed from mangrove clearing studies in the Caribbean (Lovelock et al., 2011) and Africa (Lang'at et al., 2014). These processes may be implied when we observed a decrease of dead wood carbon stocks after 5 to 10 years, following harvesting (Figure 5a). In summary, tree removal activities generate larger carbon stock losses within the biomass carbon pool rather than the soil carbon pool, despite uncertainty on the amount of direct GHG emissions.

Unlike forest harvesting, mangrove conversion into aquaculture generates carbon stocks loss from all carbon pools (Figure 5). While carbon stock losses due to harvesting can be attributed mainly to loss from the biomass pool, aquaculture conversion reduces 85% of carbon stocks from the biomass carbon pool and ~60% of carbon stocks from soil (0–50 cm) carbon pools, similar to losses observed in the Dominican Republic (Kauffman et al., 2014). In addition, further soil carbon stock losses during aquaculture development are potentially unaccounted for.

Specifically, fishpond development requires the removal of the top meter of sediments following mangrove forest clearing (Sidik et al., 2013). Therefore, mangrove conversion to other land uses, such as aquaculture, generates up to three times larger carbon stock losses compared to forest harvesting.

Our assessment reveals strong variation in mangrove carbon stocks between hydrogeomorphic settings, and therefore the degree of impact on carbon stocks and thus CO<sub>2</sub> emission may be dependent on the specific mangrove setting in which human disturbance occur. For instance, carbon stock loss caused by the same land-use change (e.g., aquaculture) may have greater impacts if aquaculture is developed over estuarine interior mangroves rather than mangroves in fringe or open coast setting. Quantifying carbon emissions from land-use change using the stock difference approach therefore requires careful selection of sampling sites, particularly pairing reference and treatment sites within a similar mangrove site and hydrogeomorphic setting.

### **Carbon stocks recovery following mangrove regeneration**

While there is no obvious change of soil carbon following harvesting, the differences in carbon stocks between some soil layers may be attributed to natural spatial variation. The carbon stocks recovery, however, was indicated by the increase of biomass carbon stocks following all regeneration years across the chronosequence (Figure 5a). The findings suggest that mangrove regeneration in the study area is more rapid compared to the global average of ~40 years (Sasmito et al., 2019) and site-specific studies in Peninsular Malaysia (> 40 years, Adame et al., 2018) and The Philippines (~50 years, Salmo et al., 2013). The efficiency of carbon stock recovery could be driven by several factors: 1) climatic conditions, as Bintuni is located near the equator with high annual rainfall (>3,000 mm), which may drive higher productivity, reduce the occurrence of natural disturbances such as cyclones, and limit variation of annual temperature and humidity; 2) hydrogeomorphic setting, as the site is located in an estuarine geomorphic setting, characterized by nutrient-rich sediment and tall forest stands (Rovai et al., 2018; Simard et al., 2019) and; 3) forest harvesting methodology, under which selective harvesting is applied in this study site (Sillanpää et al., 2017) rather than large-scale biomass removal.

## **Implications for blue carbon policy and restoration**

Indonesia's mangroves have previously been identified as a key contributor to its national carbon emissions, and a key solution to emissions reduction (Murdiyarso et al., 2015). Accounting for ~10% of the world's mangrove area, Papuan mangroves can be an important aspect of nature-based climate change mitigation in Indonesia due to their high carbon stocks, and as such they have a role to play in Indonesia's Nationally Determined Contributions (NDCs) to the Paris Agreement (Howard et al., 2017; Roe et al., 2019). In Indonesia's latest NDC, potential emissions and reductions from mangrove conversion and restoration are not separately included in accounts, but are amalgamated with other mineral soil forests under the forestry category (Government of Indonesia, 2017). Given the magnitude of soil carbon emissions from mangrove loss, and the role of mangroves in GHG management, policy makers have an opportunity to separate mangroves from other forests and account for their unique emissions profiles and removals ability, similarly to current calculations made for biogenic peatlands. When carbon emissions associated with mangrove land-use change are not separately calculated, the magnitude of this ecosystems' impact on GHG emission management may be underestimated. While previous blue carbon knowledge gaps in Indonesia were associated with the availability of suitable emissions factor data (Murdiyarso et al., 2018), findings and data from this study, along with other studies from other islands such as Kalimantan (Arifanti et al., 2019) and Sulawesi (Cameron et al., 2019), could be used by policy makers to develop science-based policy and manage blue carbon emissions abatement at the national scale.

The influence of land-use change on carbon stocks shown in this study suggests that reducing disturbance to Papua's mangroves would be an important strategy to reduce Indonesia's carbon emissions from the land use sector. Mangrove conservation would sustain the natural functions of mangroves as carbon sinks and minimize emissions from future land-use change. Papua's unprotected mangroves may face threats from proposed agricultural developments in the future (Richards & Friess, 2016). Therefore, policies should be developed to increase protected area coverage and prevent further mangrove conversion to other land uses. Such actions could

contribute efficiently to Indonesia's GHG emissions reduction targets, because the carbon loss per area and associated emissions of mangrove deforestation are between two and five times larger—depending on hydrogeomorphic setting—than emissions generated by terrestrial tropical deforestation.

This study shows that mangrove restoration, if conducted at an adequate scale, has the potential to contribute to Indonesia's NDCs by increasing mangrove carbon stocks and offsetting anthropogenic GHG emissions. This has also been shown at the site-scale for other parts of Indonesia (Cameron et al., 2019). There is growing interest in utilizing carbon removals by mangroves in Indonesia and elsewhere to finance restoration activities, by trading carbon credits through the Kyoto Protocol's Clean Development Mechanism (CDM) or voluntary Payments for Ecosystem Services (PES) schemes (Locatelli et al., 2014). However, there are clear constraints associated with the success rates of mangrove restoration (Kodikara et al., 2017) and their costs (Bayraktarov et al. 2016). Current mangrove restoration programs place a lot of attention on the low success rates of planted seedlings, because planting is conducted in inappropriate habitats adjacent to mangroves (e.g., mudflats, beaches) without regard for their hydrogeomorphic suitability (Lee et al., 2019). This study shows that the effectiveness of carbon stock recovery following mangrove regeneration is dependent on biophysical factors such as the coastal hydrogeomorphic setting. However, mangrove restoration projects are often forced into unsuitable locations due to factors such as land-use management, land tenure and inappropriate planting incentives, and these remain major constraints to successful mangrove restoration (Lovelock & Brown, 2019; Wodehouse & Rayment, 2019). Therefore, both land management and biophysical data should be equally incorporated for effective mangrove restoration to recover natural mangrove functions efficiently.

## ACKNOWLEDGMENTS

This study was a collaborative work under the Center for International Forestry Research (CIFOR)'s Sustainable Wetlands Adaptation and Mitigation Program (SWAMP) and Conservation



International Blue Carbon Initiative. We thank KESBANGPOL at The Ministry of Home Affairs for the research permit. SDS acknowledges the Australian Government RTP Scholarship and CIFOR-SWAMP Fellowship. TW was supported by the International Climate Initiative (IKI), funded by The German Federal Ministry for the Environment, Nature Conservation and Nuclear Safety (BMU). We acknowledge the fieldwork contribution of our best colleague Herry Kopalit, who passed away before the completion of the paper. We also acknowledge the fieldwork contribution from Marthinus Rumbiak, field assistants at PT BUMWI as well as CI Kaimana Office. We thank the anonymous reviewers for their constructive comments to the manuscript of this paper.

### **AUTHOR CONTRIBUTIONS:**

SDS led the study design, drafting of all figures and tables and writing of the paper with substantial contributions from DAF, LH, DM, TW, CEL, MAH, JH and MS. SDS, MS, MAH, SB, MFS, FS, BH, WYM, VIS, HH, ST, S, JDN, TP, NW, B, VN, RNY, CH and JH performed fieldwork. MS and MH contributed unpublished datasets, and TW and PZE contributed the map of mangrove hydrogeomorphic settings. SDS, MAH, JSR, and HH analyzed and interpreted data; SDS performed statistical analyses.

### **DATA SHARING AND ACCESSIBILITY**

All raw data from this study are accessible through the CIFOR Dataverse digital repository (Sasmito et al., 2019b, 2019c, 2019d).

### **REFERENCES**

- Adame, M.F., Kauffman, J.B., Medina, I., Gamboa, J.N., Torres, O., Caamal, J.P., Reza, M. & Herrera-Silveira, J.A. (2013). Carbon stocks of tropical coastal wetlands within the karstic landscape of the Mexican Caribbean. *PLoS One*, **8**, e56569.
- Adame, M. F., Cherian, S., Reef, R., & Stewart-Koster, B. (2017). Mangrove root biomass and the uncertainty of belowground carbon estimations. *Forest ecology and management*, *403*, 52-60.

Adame, M. F., Zakaria, R. M., Fry, B., Chong, V. C., Then, Y. H. A., Brown, C. J., & Lee, S. Y. (2018). Loss and recovery of carbon and nitrogen after mangrove clearing. *Ocean & Coastal Management*, 161, 117-126. doi:10.1016/j.ocecoaman.2018.04.019

Arifanti, V.B., Kauffman, J.B., Hadriyanto, D., Murdiyarso, D. & Diana, R. (2019). Carbon dynamics and land use carbon footprints in mangrove-converted aquaculture: The case of the Mahakam Delta, Indonesia. *Forest Ecology and Management*, **432**, 17-29.

Atwood, T. B., Connolly, R. M., Almahasheer, H., Carnell, P. E., Duarte, C. M., Ewers Lewis, C. J., . . . Lovelock, C. E. (2017). Global patterns in mangrove soil carbon stocks and losses. *Nature Climate Change*, **7**(7), 523-528. doi:10.1038/nclimate3326

Bayraktarov, E., Saunders, M. I., Abdullah, S., Mills, M., Beher, J., Possingham, H. P., ... & Lovelock, C. E. (2016). The cost and feasibility of marine coastal restoration. *Ecological Applications*, **26**(4), 1055-1074.

Bouillon, S., Borges, A.V., Castañeda-Moya, E., Diele, K., Dittmar, T., Duke, N.C., Kristensen, E., Lee, S.Y., Marchand, C., Middelburg, J.J., Rivera-Monroy, V.H., Smith, T.J. & Twilley, R.R. (2008). Mangrove production and carbon sinks: A revision of global budget estimates. *Global Biogeochemical Cycles*, **22**.

Cameron, C., Hutley, L. B., Friess, D. A., & Brown, B. (2019). Community structure dynamics and carbon stock change of rehabilitated mangrove forests in Sulawesi, Indonesia. *Ecological applications*, **29**(1), e01810.

Donato, D.C., Kauffman, J.B., Murdiyarso, D., Kurnianto, S., Stidham, M. & Kanninen, M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, **4**, 293-297.

Ellert, B., Janzen, H., VandenBygaart, A., & Bremer, E. (2007). Measuring Change in Soil Organic Carbon Storage. *Soil sampling and methods of analysis*, 25.

Hamilton, S.E. & Casey, D. (2016). Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC-21). *Global Ecology and Biogeography*, **25**, 729-738.

Government of Indonesia. (2017). Nationally determined contribution (NDC) and its progress of implementation. 54pp. Retrieved from [http://ditjenppi.menlhk.go.id/reddplus/images/adminppi/dokumen/summary\\_NDC\\_english\\_opt.pdf](http://ditjenppi.menlhk.go.id/reddplus/images/adminppi/dokumen/summary_NDC_english_opt.pdf)

Hamilton, S.E. & Friess, D.A. (2018). Global carbon stocks and potential emissions due to mangrove deforestation from 2000 to 2012. *Nature Climate Change*, **8**, 240.

- Howard, J., Hoyt, S., Isensee, K., Pidgeon, E., Telszewski, M. (2014). Coastal Blue Carbon: Methods for assessing carbon stocks and emissions factors in mangroves, tidal salt marshes, and seagrass meadows. Conservation International, Intergovernmental Oceanographic Commission of UNESCO, International Union for Conservation of Nature. Arlington, Virginia, USA.
- Howard, J., Sutton-Grier, A., Herr, D., Kleyapas, J., Landis, E., McLeod, E., . . . Simpson, S. (2017). Clarifying the role of coastal and marine systems in climate mitigation. *Frontiers in Ecology and the Environment*, 15(1), 42-50. doi:10.1002/fee.1451
- Kauffman, J. & Donato, D. (2012). Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in mangrove forests. *CIFOR Working Paper* No. 86. Bogor, Indonesia.
- Kauffman, J.B., Heider, C., Norfolk, J. & Payton, F. (2014). Carbon stocks of intact mangroves and carbon emissions arising from their conversion in the Dominican Republic. *Ecological Applications*, 24, 518-527.
- Kauffman, J.B., Heider, C., Cole, T.G., Dwire, K.A. & Donato, D.C. (2011). Ecosystem Carbon Stocks of Micronesian Mangrove Forests. *Wetlands*, 31, 343-352.
- Kauffman, J.B., Bernardino, A.F., Ferreira, T.O., Bolton, N.W., Gomes, L.E.O. & Nobrega, G.N. (2018). Shrimp ponds lead to massive loss of soil carbon and greenhouse gas emissions in northeastern Brazilian mangroves. *Ecol Evol*, 8, 5530-5540.
- Kauffman, J.B., M.F. Adame, V.B. Arifanti, L.M. Schile-Beers, A.F. Bernardino, R.K. Bhomia, D.C. Donato, I. Feller, T.O. Ferreira, M.C.J. Garcia, R.A. MacKenzie, J.P. Megonigal, D. Murdiyarso, L. Simpson, and H.H. Trejo. (2020). Total ecosystem carbon stocks of mangroves across broad global environmental and physical gradients. *Ecological Monographs*. doi: <https://doi.org/10.1002/ecm.1405>
- Kauffman, J.B., Arifanti, V.B., Hernández Trejo, H., del Carmen Jesús García, M., Norfolk, J., Cifuentes, M., Hadriyanto, D. & Murdiyarso, D. (2017). The jumbo carbon footprint of a shrimp: carbon losses from mangrove deforestation. *Frontiers in Ecology and the Environment*, 15, 183-188.
- Kodikara, K. A. S., Mukherjee, N., Jayatissa, L. P., Dahdouh-Guebas, F., & Koedam, N. (2017). Have mangrove restoration projects worked? An in-depth study in Sri Lanka. *Restoration Ecology*, 25(5), 705-716.
- Krauss, K. W., McKee, K. L., Lovelock, C. E., Cahoon, D. R., Saintilan, N., Reef, R., & Chen, L. (2014). How mangrove forests adjust to rising sea level. *New Phytol*, 202(1), 19-34. doi:10.1111/nph.12605
- Kusmana, C., & Onrizal, S. (2003). Jenis–Jenis Pohon Mangrove di Teluk Bintuni. Bogor (ID): Fakultas Kehutanan IPB dan PT. Bintuni Utama Murni Wood Industries.

- Lang'at, J. K., Kairo, J. G., Mencuccini, M., Bouillon, S., Skov, M. W., Waldron, S., & Huxham, M. (2014). Rapid losses of surface elevation following tree girdling and cutting in tropical mangroves. *PLoS One*, 9(9), e107868. doi:10.1371/journal.pone.0107868
- Lee, S. Y., Hamilton, S., Barbier, E. B., Primavera, J., & Lewis III, R. R. (2019). Better restoration policies are needed to conserve mangrove ecosystems. *Nat Ecol Evol*. doi:10.1038/s41559-019-0861-y
- Locatelli, T., Binet, T., Kairo, J. G., King, L., Madden, S., Patenaude, G., ... & Huxham, M. (2014). Turning the tide: how blue carbon and payments for ecosystem services (PES) might help save mangrove forests. *Ambio*, 43(8), 981-995.
- Lovelock, C.E. & Duarte, C.M. (2019). Dimensions of Blue Carbon and emerging perspectives. *Biol Lett*, **15**, 20180781.
- Lovelock, C. E., Ruess, R. W., & Feller, I. C. (2011). CO<sub>2</sub> efflux from cleared mangrove peat. *PLoS One*, 6(6), e21279. doi:10.1371/journal.pone.0021279
- Lovelock, C. E., & Brown, B. M. (2019). Land tenure considerations are key to successful mangrove restoration. *Nature Ecology & Evolution*, 3(8), 1135-1135. doi:10.1038/s41559-019-0942-y
- Murdiyarso, D., Purbopuspito, J., Kauffman, J.B., Warren, M.W., Sasmito, S.D., Donato, D.C., Manuri, S., Krisnawati, H., Taberima, S. & Kurnianto, S. (2015). The potential of Indonesian mangrove forests for global climate change mitigation. *Nature Climate Change*, **5**, 1089-1092.
- Murdiyarso, D., Sukara, E., Supriatna, J., Koropitan, A., Juliandi, B., Jompa, J. (2018). Creating blue carbon opportunities in the maritime archipelago Indonesia. *CIFOR Policy Brief* No 3. Bogor, Indonesia. doi: 10.17528/cifor/007058
- Nam, V.N., Sasmito, S.D., Murdiyarso, D., Purbopuspito, J. & MacKenzie, R.A. (2016). Carbon stocks in artificially and naturally regenerated mangrove ecosystems in the Mekong Delta. *Wetlands Ecology and Management*, **24**, 231-244.
- Paz, C. P., Goosem, M., Bird, M., Preece, N., Goosem, S., Fensham, R., & Laurance, S. (2016). Soil types influence predictions of soil carbon stock recovery in tropical secondary forests. *Forest Ecology and Management*, 376, 74-83.
- Richards, D.R. & Friess, D.A. (2016). Rates and drivers of mangrove deforestation in Southeast Asia, 2000-2012. *Proc Natl Acad Sci U S A*, **113**, 344-9.
- Roe, S., Streck, C., Obersteiner, M., Frank, S., Griscom, B., Drouet, L., . . . Lawrence, D. (2019). Contribution of the land sector to a 1.5 °C world. *Nature Climate Change*. doi:10.1038/s41558-019-0591-9
- Rogers, K., Kelleway, J. J., Saintilan, N., Megonigal, J. P., Adams, J. B., Holmquist, J. R., . . . Woodroffe, C. D. (2019). Wetland carbon storage controlled by millennial-scale variation in relative sea-level rise. *Nature*, 567(7746), 91-95. doi:10.1038/s41586-019-0951-7

Rovai, A. S., Twilley, R. R., Castañeda-Moya, E., Riul, P., Cifuentes-Jara, M., Manrow-Villalobos, M., . . .

Pagliosa, P. R. (2018). Global controls on carbon storage in mangrove soils. *Nature Climate Change*, 8(6), 534-538. doi:10.1038/s41558-018-0162-5

Sasmito, S.D., Kuzyakov, Y., Lubis, A.A., Murdiyarso, D., Hutley, L.B., Bachri, S., Friess, D.A., Martius, C.M., Borchard, N. 2020. Carbon burial rates and sources in soils of coastal mudflat and mangrove ecosystems. *Catena*, 184. doi: 10.1016/j.catena.2019.104414

Sasmito SD, Taillardat P, Clendenning JN, Cameron C, Friess DA, Murdiyarso D, Hutley LB. (2019a). Effect of land-use and land-cover change on mangrove blue carbon: A systematic review. *Global Change Biology*, 25(12), 4291-4302. doi: 10.1111/gcb.14774

Sasmito SD, Sillanpää M, Hayes MA, Bachri S, Saragi-Sasmito MF, Sidik F, Hanggara B, Mofu WY, Rumbiak VI, Hendri, Rumbiak MI, Taberima S, Suhaemi, Nugroho JD, Pattiasina TF, Widagti N, Barakalla, Rahajoe JS, Hartantri H, Nikijuluw V, Jowey RN, Heatunubun C, Ermgassen PZ, Worthington TA, Howard J, Lovelock CE, Friess DA, Hutley LB, Murdiyarso D. (2019b). SWAMP Dataset-Mangrove biomass vegetation-West Papua-2019. Center for International Forestry Research (CIFOR), V1. <https://doi.org/10.17528/CIFOR/DATA.00193>

Sasmito SD, Sillanpää M, Hayes MA, Bachri S, Saragi-Sasmito MF, Sidik F, Hanggara B, Mofu WY, Rumbiak VI, Hendri, Rumbiak MI, Taberima S, Suhaemi, Nugroho JD, Pattiasina TF, Widagti N, Barakalla, Rahajoe JS, Hartantri H, Nikijuluw V, Jowey RN, Heatunubun C, Ermgassen PZ, Worthington TA, Howard J, Lovelock CE, Friess DA, Hutley LB, Murdiyarso D. (2019c). SWAMP Dataset-Mangrove soil carbon-West Papua-2019. Center for International Forestry Research (CIFOR), V1. <https://doi.org/10.17528/CIFOR/DATA.00192>

Sasmito SD, Sillanpää M, Hayes MA, Bachri S, Saragi-Sasmito MF, Sidik F, Hanggara B, Mofu WY, Rumbiak VI, Hendri, Rumbiak MI, Taberima S, Suhaemi, Nugroho JD, Pattiasina TF, Widagti N, Barakalla, Rahajoe JS, Hartantri H, Nikijuluw V, Jowey RN, Heatunubun C, Ermgassen PZ, Worthington TA, Howard J, Lovelock CE, Friess DA, Hutley LB, Murdiyarso D. (2019d). SWAMP Dataset-Mangrove necromass-West Papua-2019. Center for International Forestry Research (CIFOR), V1. <https://doi.org/10.17528/CIFOR/DATA.00194>

Salmo, S. G., Lovelock, C., & Duke, N. C. (2013). Vegetation and soil characteristics as indicators of restoration trajectories in restored mangroves. *Hydrobiologia*, 720(1), 1-18. doi:10.1007/s10750-013-1617-3

Sidik, F., & Lovelock, C. E. (2013). CO<sub>2</sub> efflux from shrimp ponds in Indonesia. *PLoS One*, 8(6), e66329. doi:10.1371/journal.pone.0066329


- Sillanpää, M., Vantellingen, J. & Friess, D.A. (2017). Vegetation regeneration in a sustainably harvested mangrove forest in West Papua, Indonesia. *Forest Ecology and Management*, **390**, 137-146.
- Simard, M., Fatoyinbo, L., Smetanka, C., Rivera-Monroy, V.H., Castañeda-Moya, E., Thomas, N. & Van der Stocken, T. (2019). Mangrove canopy height globally related to precipitation, temperature and cyclone frequency. *Nature Geoscience*, **12**, 40-45.
- Stringer, C.E., Trettin, C.C., Zarnoch, S.J. & Tang, W. (2015). Carbon stocks of mangroves within the Zambezi River Delta, Mozambique. *Forest Ecology and Management*, **354**, 139-148.
- Taillardat, P., Friess, D. A., & Lupascu, M. (2018). Mangrove blue carbon strategies for climate change mitigation are most effective at the national scale. *Biol Lett*, *14*(10). doi:10.1098/rsbl.2018.0251
- Twilley, R. R., Rovai, A. S., & Riul, P. (2018). Coastal morphology explains global blue carbon distributions. *Frontiers in Ecology and the Environment*. doi:10.1002/fee.1937
- Wodehouse, D. C., & Rayment, M. B. (2019). Mangrove area and propagule number planting targets produce sub-optimal rehabilitation and afforestation outcomes. *Estuarine, Coastal and Shelf Science*, *222*, 91-102.
- Woodroffe, C. D., Rogers, K., McKee, K. L., Lovelock, C. E., Mendelssohn, I. A., & Saintilan, N. (2016). Mangrove Sedimentation and Response to Relative Sea-Level Rise. *Ann Rev Mar Sci*, *8*, 243-266. doi:10.1146/annurev-marine-122414-034025
- Worthington, T., & Spalding, M. (2018). Mangrove Restoration Potential: A global map highlighting a critical opportunity. <https://doi.org/10.17863/CAM.39153>

Table 1. Summary of mangrove settings, land-use types, year of field sampling and number of sampled plots across study sites.


Site	Macro-scale setting (hydrogeomorphic variation)	Meso-scale setting (hydrodynamic or tidal elevation)	Land-use description	Sampling year	Number of plots	Soil depth (cm)	Dominant species
Arguni Bay	Estuarine	Interior	Undisturbed forest	2015	41	300	<i>Rhizophora</i> spp. (41%)
Bintuni Bay	Estuarine	Interior	Undisturbed forest	2018	18	300	<i>Rhizophora</i> <i>apiculata</i> (33%)
	Open coast	Interior	Harvested forest	2018	18	300	—
	Estuarine	Interior	Post-harvest forest (5 years old)	2018	18	300	<i>Rhizophora</i> <i>apiculata</i> (44%)
	Estuarine	Interior	Post-harvest forest (10 years old)	2018	18	286	<i>Rhizophora</i> <i>apiculata</i> (91%)
	Estuarine	Interior	Post-harvest forest (15 years old)	2018	18	300	<i>Rhizophora</i> <i>apiculata</i> (88%)
	Estuarine	Interior	Post-harvest forest (25 years old)	2018	18	300	<i>Rhizophora</i> <i>apiculata</i>

							(81%)
Buruway	Estuarine	Interior	Undisturbed forest	2017	17	230	<i>Rhizophora apiculata</i> (37%)
	Estuarine	Fringe	Undisturbed forest	2017	28	107	<i>Rhizophora apiculata</i> (36%)
Etna	Open coast	Interior	Undisturbed forest	2017	43	234	<i>Rhizophora apiculata</i> (47%)
	Open coast	Fringe	Undisturbed forest	2017	6	190	<i>Rhizophora apiculata</i> (50%)
Kaimana	Estuarine	Interior	Undisturbed forest	2017	5	113	<i>Rhizophora mucronata</i> (39%)
	Estuarine	Interior	Aquaculture (15 years old)	2017	7	56	<i>Rhizophora apiculata</i> (49%)
Total					255		







gcb\_15056\_f1.jpg




gcb\_15056\_f2.jpg



# All undisturbed mangrove soil samples (n=718)



# Bintuni Bay mangrove soil samples (n=324)



gcb\_15056\_f4.jpg

